

# The EPEC Model

R. Fitzpatrick\*  
Institute for Fusion Studies,  
Department of Physics,  
University of Texas at Austin,  
Austin TX 78712, USA

September 28, 2021

## 1 Plasma Response in Outer Region

### 1.1 Coordinates

Let  $R$ ,  $\phi$ , and  $Z$  be right-handed cylindrical coordinates whose symmetry axis corresponds to the toroidal symmetry axis of the plasma. Let  $r$ ,  $\theta$ , and  $\phi$  be right-handed flux coordinates whose Jacobian is  $\mathcal{J} \equiv (\nabla r \times \nabla \theta \cdot \nabla \phi)^{-1} = r R^2 / R_0$ . Here,  $R_0$  is a convenient scale major radius,  $r$  is a magnetic flux-surface label with dimensions of length, and  $\theta$  is an axisymmetric angular coordinate that increases by  $2\pi$  radians for every poloidal circuit of the magnetic axis. Let  $r = 0$  correspond to the magnetic axis, and let  $r = r_{100}$  correspond to the LCFS. Let  $\theta = 0$  correspond to the inboard midplane, and let  $0 < \theta < \pi$  correspond to the region above the midplane.

### 1.2 Equilibrium Magnetic Field

The equilibrium magnetic field is written  $\mathbf{B} = R_0 B_0 [f(r) \nabla \phi \times \nabla r + g(r) \nabla \phi]$ , where  $B_0$  is a convenient scale toroidal magnetic field-strength, and  $q(r) = r g / (R_0 f)$  is the safety-factor profile. The equilibrium poloidal magnetic flux (divided by  $2\pi$ ),  $\Psi_p(r)$ , satisfies  $d\Psi_p/dr = R_0 B_0 f(r)$ , where, by convention,  $\Psi_p(r_{100}) = 0$ . The normalized poloidal magnetic flux,  $\Psi_N(r)$ , is defined such that  $\Psi_N(r) = 1 - \Psi_p(r)/\Psi_p(0)$ . Hence,  $\Psi_N(0) = 0$  and  $\Psi_N(r_{100}) = 1$ . Finally, if  $\Psi_N(r_{95}) = 0.95$  then  $q_{95} \equiv q(r_{95})$ .

---

\*rfitzp@utexas.edu

### 1.3 Perturbed Magnetic Field

Consider the response of the plasma to an RMP with  $n > 0$  periods in the toroidal direction. We can write the components of the perturbed magnetic field in the form

$$\frac{r R^2 \delta \mathbf{B} \cdot \nabla r}{R_0^2} = i \sum_j \psi_j(r) e^{i(m_j \theta - n \phi)}, \quad (1)$$

$$\mathcal{J} \delta \mathbf{B} \cdot \nabla \phi \times \nabla r = - \sum_j \Xi_j(r) e^{i(m_j \theta - n \phi)}, \quad (2)$$

$$R^2 \delta \mathbf{B} \cdot \nabla \phi = n \sum_j \frac{\bar{\Xi}_j(r)}{m_j} e^{i(m_j \theta - n \phi)}, \quad (3)$$

where the sum is over all relevant poloidal harmonics of the perturbed magnetic field.

Let there be  $K$  resonant magnetic flux-surfaces in the plasma, labelled 1 through  $K$ . Consider the  $k$ th resonant surface,  $r = r_k$ , at which  $n q(r_k) = m_k$ , where  $m_k$  is a positive integer. Let  $\Psi_k = \psi_k(r_k)/m_k$ , and  $\Delta \Psi_k = [\Xi_k]_{r_k^-}^{r_k^+}$ . Here,  $\Psi_k$  is the (complex) reconnected helical magnetic flux (divided by  $2\pi R_0$ ) at the  $k$ th resonant surface, whereas  $\Delta \Psi_k$  (which has the same dimensions as  $\Psi_k$ ) is a (complex) measure of the strength of the current sheet (consisting of filaments running parallel to the local equilibrium magnetic field) at the same resonant surface.

### 1.4 Toroidal Tearing Mode Dispersion Relation

In the presence of the RMP, the  $\Psi_k$  and the  $\Delta \Psi_k$  values are related according to the inhomogeneous toroidal tearing mode dispersion relation, which takes the form

$$\Delta \Psi_k = \sum_{k'=1, K} E_{kk'} \Psi_{k'} + |E_{kk}| \chi_k. \quad (4)$$

Here,  $E_{kk'}$  (for  $k, k' = 1, K$ ) is the dimensionless, Hermitian, toroidal tearing mode stability matrix, whereas the  $\chi_k$  (for  $k = 1, K$ ) parameterize the current sheets driven at the various resonant surfaces when the plasma responds to the applied RMP in accordance with the equations of linearized, marginally-stable, ideal-MHD.

The EPEC model determines the elements of the  $E_{kk'}$  matrix using a high- $q$  approximation. In fact, if  $F_{kk'}$  is the inverse of the  $E_{kk'}$  matrix then

$$F_{kk'} = \oint \oint G(R_k, Z_k; R_{k'}, Z_{k'}) e^{-i(m_k \theta_k - m_{k'} \theta_{k'})} \frac{d\theta_k}{2\pi} \frac{d\theta_{k'}}{2\pi}, \quad (5)$$

and

$$G(R_k, Z_k; R_{k'}, Z_{k'}) = \frac{(-1)^n \pi^2 R_k R_{k'} / R_0}{2 \Gamma(1/2) \Gamma(n + 1/2)} \left[ \frac{\cosh \eta_{kk'}}{R_k^2 + R_{k'}^2 + (Z_k - Z_{k'})^2} \right]^{1/2}$$

$$\times \left[ (n - 1/2) P_{-1/2}^{n-1}(\cosh \eta_{kk'}) + \frac{P_{-1/2}^{n+1}(\cosh \eta_{kk'})}{n + 1/2} \right], \quad (6)$$

with

$$\eta_{kk'} = \tanh^{-1} \left[ \frac{2 R_k R_{k'}}{R_k^2 + R_{k'}^2 + (Z_k - Z_{k'})^2} \right]. \quad (7)$$

Here, the double integral in Eq. (5) is taken around the  $k$ th resonant surface (cylindrical coordinates  $R_k, 0, Z_k$ ; flux coordinates  $r_k, \theta_k, 0$ , with  $r_k$  constant; resonant poloidal mode number  $m_k$ ) and the  $k'$ th resonant surface (cylindrical coordinates  $R_{k'}, 0, Z_{k'}$ ; flux coordinates  $r_{k'}, \theta_{k'}, 0$ , with  $r_{k'}$  constant; resonant poloidal mode number  $m_{k'}$ ). Finally, the  $\Gamma(z)$  and  $P_\mu^\nu(z)$  are gamma functions and associated Legendre functions, respectively.

The (complex)  $\chi_k$  parameters are determined from the GPEC code. To be more exact, the GPEC code calculates the (complex) dimensionless  $\Delta_{m_k n}$  parameters which measure the strengths of the ideal current sheets that develop at the various resonant magnetic flux-surfaces in the plasma in response to the applied RMP. The  $\Delta_{m_k n}$  parameters are related to the  $\chi_k$  parameters according to

$$\frac{\chi_k}{R_0 B_0} = -i \frac{\Delta_{m_k n}}{|E_{kk}|} \left( \frac{r_k}{R_0} \right)^2 \frac{g(r_k)}{m_k [a_{kk}(r_k) + (r_k/R_0 q_k)^2]}, \quad (8)$$

where  $q_k = m_k/n$ , and  $a_{kk}(r) = \oint |\nabla r|^{-2} d\theta / (2\pi)$ .

## 1.5 Glasser-Greene-Johnson Data

Let us write

$$\mathbf{B} = \nabla\phi \times \nabla\psi_p + g(\psi_p) \nabla\phi. \quad (9)$$

The angle  $\theta^*$  is defined such that

$$\nabla\phi \times \nabla\psi_p \cdot \nabla\theta^* = \gamma(\psi_p) B. \quad (10)$$

Let

$$J_1(\psi_p) = \oint \frac{1}{B} \frac{d\theta^*}{2\pi}, \quad (11)$$

$$J_2(\psi_p) = \oint B \frac{d\theta^*}{2\pi}, \quad (12)$$

$$J_3(\psi_p) = \oint \frac{1}{B^3} \frac{d\theta^*}{2\pi}, \quad (13)$$

$$J_4(\psi_p) = \oint \frac{1}{B |\nabla\psi_p|^2} \frac{d\theta^*}{2\pi}, \quad (14)$$

$$J_5(\psi_p) = \oint \frac{B}{|\nabla\psi_p|^2} \frac{d\theta^*}{2\pi}, \quad (15)$$

$$J_6(\psi_p) = \oint \frac{1}{B^3 |\nabla\psi_p|^2} \frac{d\theta^*}{2\pi}. \quad (16)$$

It follows that

$$E(\psi_p) = -\frac{dP/d\psi_p}{(dq/d\psi_p)^2} \frac{1}{\gamma} \left[ \frac{d}{d\psi_p} \left( \frac{J_1}{\gamma} \right) - g \frac{dq}{d\psi_p} \frac{J_1}{J_2} \right] J_5, \quad (17)$$

$$F(\psi_p) = \frac{(dP/d\psi_p)^2}{(dq/d\psi_p)^2} \frac{1}{\gamma^2} [g^2 (J_5 J_6 - J_4^2) + J_5 J_3], \quad (18)$$

$$H(\psi_p) = \frac{dP/d\psi_p}{dq/d\psi_p} \frac{g}{\gamma} \left( J_4 - \frac{J_1 J_5}{J_2} \right), \quad (19)$$

where  $P(\psi_p)$  is the total plasma pressure. Finally,

$$D_R(\psi_p) = E + F + H^2. \quad (20)$$

## 1.6 Cylindrical Tearing Mode Equation

The cylindrical tearing mode equation for the eigenfunction,  $\psi(r)$ , of the  $m_k, n$  tearing mode takes the form

$$\frac{d^2\psi}{dr^2} + \frac{1}{r} \frac{d\psi}{dr} - \frac{m_k^2}{r^2} \psi - \frac{\hat{J}'_\phi \psi}{r(1/q - n/m_k)} = 0. \quad (21)$$

Here,

$$\hat{J}'_\phi(r) = -\frac{\mu_0 R_0 R}{B_0} \frac{dP}{d\Psi_p} - R_0^2 B_0 g \frac{dg}{d\Psi_p}, \quad (22)$$

where  $P(\Psi_p)$  is the total plasma pressure, the previous expression is evaluated on the out-board mid-plane, and  $\hat{J}'_\phi \equiv d\hat{J}_\phi/dr$ .

Suppose that the plasma is surrounded by a thin resistive wall of time constant  $\tau_w$ , located at  $r = r_w > r_{100}$ . The quantities  $\Delta_k^{nw}$  and  $\Sigma_k^{nw}$  are defined as

$$\Delta_k^{nw} = \lim_{\delta \rightarrow 0} \left[ \frac{r \psi'_{nw}}{\psi_{nw}} \Big|_{r_k + \delta} - \frac{r \psi'_{nw}}{\psi_{nw}} \Big|_{r_k - \delta} \right], \quad (23)$$

$$\Sigma_k^{nw} = \lim_{\delta \rightarrow 0} \left[ \frac{r \psi'_{nw}}{\psi_{nw}} \Big|_{r_k + \delta} + \frac{r \psi'_{nw}}{\psi_{nw}} \Big|_{r_k - \delta} - 2 A_k (1 + \ln \delta) \right], \quad (24)$$

where

$$A_k = - \left( \frac{r q \hat{J}'_\phi}{s} \right)_{r_k}, \quad (25)$$

$$s(r) = \frac{d \ln q}{d \ln r}. \quad (26)$$

Here,  $\psi_{nw}(r)$  is a solution of the cylindrical tearing mode equation that is well-behaved at  $r = 0$ , continuous at  $r = r_k$ , and satisfies

$$\left. \frac{r \psi'_{nw}}{\psi_{nw}} \right|_{r_{100}} = -m. \quad (27)$$

Moreover,  $\psi' \equiv d\psi/dr$ .

The quantities  $\Delta_k^{pw}$  and  $\Sigma_k^{pw}$  are defined as

$$\Delta_k^{nw} = \lim_{\delta \rightarrow 0} \left[ \left. \frac{r \psi'_{pw}}{\psi_{pw}} \right|_{r_k + \delta} - \left. \frac{r \psi'_{pw}}{\psi_{pw}} \right|_{r_k - \delta} \right], \quad (28)$$

$$\Sigma_k^{pw} = \lim_{\delta \rightarrow 0} \left[ \left. \frac{r \psi'_{pw}}{\psi_{pw}} \right|_{r_k + \delta} + \left. \frac{r \psi'_{pw}}{\psi_{pw}} \right|_{r_k - \delta} - 2 A_k (1 + \ln \delta) \right]. \quad (29)$$

Here,  $\psi_{pw}(r)$  is a solution of the cylindrical tearing mode equation that is well-behaved at  $r = 0$ , continuous at  $r = r_k$ , and satisfies

$$\left. \frac{r \psi'_{pw}}{\psi_{pw}} \right|_{r_{100}} = -m \left[ \frac{1 + (r_{100}/r_w)^{2m}}{1 - (r_{100}/r_w)^{2m}} \right]. \quad (30)$$

It is also helpful to define

$$\Sigma_{kw} = \frac{2m (r_{100}/r_w)^m}{1 - (r_{100}/r_w)^{2m}} \frac{\psi_{pw}(r_{100})}{\psi_{pw}(r_k)}, \quad (31)$$

$$\Delta_{kw} = - \left( \frac{\Sigma_{kw}^2}{\Delta_k^{nw} - \Delta_k^{pw}} \right), \quad (32)$$

$$B_k = - \left( \frac{r^2 q \hat{J}_\phi''}{s} \right)_{r_k}, \quad (33)$$

where  $\hat{J}_\phi'' \equiv d^2 \hat{J}_\phi / dr^2$ .

## 2 Neoclassical Physics

### 2.1 Plasma Species

The plasma is assumed to consist of three (charged) species (with Maxwellian distribution functions); namely, electrons ( $e$ ), majority ions ( $i$ ), and impurity ions ( $I$ ). The charges of the three species are  $e_e = -e$ ,  $e_i = e$ , and  $e_I = Z_I e$ , respectively, where  $e$  is the magnitude of the electron charge. Quasi-neutrality demands that  $n_e = n_i + Z_I n_I$ , where  $n_a(r)$  is the species- $a$  number density. Let  $\alpha_I(r) = Z_I (Z_{\text{eff}} - 1) / (Z_I - Z_{\text{eff}})$ , where  $Z_{\text{eff}}(r) = (n_i + Z_I^2 n_I) / n_e$  is the effective ion charge number. It follows that  $n_i/n_e = (Z_I - Z_{\text{eff}}) / (Z_I - 1)$  and  $n_I/n_e = (Z_{\text{eff}} - 1) / [Z_I (Z_I - 1)]$ . Finally, let  $Z_{\text{eff } i} = (Z_I - Z_{\text{eff}}) / (Z_I - 1)$ , and  $Z_{\text{eff } I} = Z_I (Z_{\text{eff}} - 1) / (Z_I - 1)$ . Note that  $Z_{\text{eff}} = Z_{\text{eff } i} + Z_{\text{eff } I}$ .

## 2.2 Collisionality Parameters

Consider an equilibrium magnetic flux-surface whose label is  $r$ . Let

$$\frac{1}{\gamma(r)} = \frac{q}{g} \oint \frac{B R^2}{B_0 R_0^2} \frac{d\theta}{2\pi}, \quad (34)$$

where  $B = |\mathbf{B}|$ . It is helpful to define a new poloidal angle  $\Theta$  such that

$$\frac{d\Theta}{d\theta} = \frac{\gamma q}{g} \frac{B R^2}{B_0 R_0^2}. \quad (35)$$

Let

$$I_1 = \oint \frac{B_0}{B} \frac{d\Theta}{2\pi}, \quad (36)$$

$$I_2 = \oint \frac{B}{B_0} \frac{d\Theta}{2\pi}, \quad (37)$$

$$I_3 = \oint \left( \frac{\partial B}{\partial \Theta} \right)^2 \frac{1}{B_0 B} \frac{d\Theta}{2\pi}, \quad (38)$$

$$I_{4,j} = \sqrt{2j} \oint \frac{\cos(j\Theta)}{B/B_0} \frac{d\Theta}{2\pi}, \quad (39)$$

$$I_{5,j} = \sqrt{2j} \oint \frac{\cos(j\Theta)}{2(B/B_0)^2} \frac{d\Theta}{2\pi}, \quad (40)$$

$$I_6(\lambda) = \oint \frac{\sqrt{1 - \lambda B/B_{\max}}}{B/B_0} \frac{d\Theta}{2\pi}, \quad (41)$$

where  $B_{\max}$  is the maximum value of  $B$  on the magnetic flux-surface, and  $j$  a positive integer. The species- $a$  transit frequency is written  $\omega_{ta}(r) = K_t \gamma v_{Ta}$ , where

$$K_t(r) = \frac{I_1^2 I_3}{I_2^2 \sum_{j=1,\infty} I_{4,j} I_{5,j}}, \quad (42)$$

and  $v_{Ta} = \sqrt{2T_a/m_a}$ . Here,  $m_a$  is the species- $a$  mass, and  $T_a(r)$  the species- $a$  temperature (in energy units). The fraction of circulating particles is

$$f_c(r) = \frac{3 I_2}{4} \frac{B_0^2}{B_{\max}^2} \int_0^1 \frac{\lambda d\lambda}{I_6(\lambda)}. \quad (43)$$

Finally, the dimensionless species- $a$  collisionality parameter is written  $\nu_{*a}(r) = K_* g_t / (\omega_{ta} \tau_{aa})$ , where  $g_t(r) = f_c / (1 - f_c)$ ,

$$K_*(r) = \frac{3}{8\pi} \frac{I_2}{I_3} K_t^2, \quad (44)$$

$$\frac{1}{\tau_{aa}(r)} = \frac{4}{3\sqrt{\pi}} \frac{4\pi n_a e_a^4 \ln \Lambda}{(4\pi \epsilon_0)^2 m_a^2 v_{Ta}^3}. \quad (45)$$

Here, the Coulomb logarithm,  $\ln \Lambda$ , is assumed to take the same large constant value (i.e.,  $\ln \Lambda \simeq 17$ ), independent of species.

### 2.3 Collisional Friction Matrices

Let  $x_{ab} = v_{Tb}/v_{Ta}$ . In the following, all quantities that are of order  $(m_e/m_i)^{1/2}$ ,  $(m_e/m_I)^{1/2}$ , or smaller, are neglected with respect to unity. The  $2 \times 2$  dimensionless ion collisional friction matrices,  $[F^{ii}](r)$ ,  $[F^{iI}](r)$ ,  $[F^{Ii}](r)$ , and  $[F^{II}](r)$ , are defined to have the following elements:

$$F_{00}^{ii} = \frac{\alpha_I (1 + m_i/m_I)}{(1 + x_{iI}^2)^{3/2}}, \quad (46)$$

$$F_{01}^{ii} = \frac{3}{2} \frac{\alpha_I (1 + m_i/m_I)}{(1 + x_{iI}^2)^{5/2}}, \quad (47)$$

$$F_{11}^{ii} = \sqrt{2} + \frac{\alpha_I [13/4 + 4x_{iI}^2 + (15/2)x_{iI}^4]}{(1 + x_{iI}^2)^{5/2}}, \quad (48)$$

$$F_{01}^{iI} = \frac{3}{2} \frac{T_i}{T_I} \frac{\alpha_I (1 + m_I/m_i)}{x_{iI} (1 + x_{iI}^2)^{5/2}}, \quad (49)$$

$$F_{11}^{iI} = \frac{27}{4} \frac{T_i}{T_I} \frac{\alpha_I x_{iI}^2}{(1 + x_{iI}^2)^{5/2}}, \quad (50)$$

$$F_{11}^{Ii} = \frac{27}{4} \frac{\alpha_I x_{iI}^2}{(1 + x_{iI}^2)^{5/2}}, \quad (51)$$

$$F_{11}^{II} = \frac{T_i}{T_I} \left\{ \sqrt{2} \alpha_I^2 x_{iI} + \frac{\alpha_I [15/2 + 4x_{iI}^2 + (13/4)x_{iI}^4]}{(1 + x_{iI}^2)^{5/2}} \right\}, \quad (52)$$

$$F_{10}^{ii} = F_{01}^{ii}, F_{00}^{ii} = F_{00}^{ii}, F_{10}^{iI} = F_{01}^{iI}, F_{00}^{iI} = F_{00}^{iI}, F_{01}^{iI} = F_{01}^{iI}, F_{10}^{Ii} = F_{01}^{Ii}, F_{00}^{Ii} = F_{00}^{Ii}, F_{01}^{Ii} = F_{01}^{Ii}, F_{10}^{II} = F_{00}^{II}, F_{01}^{II} = F_{01}^{II}, F_{10}^{II} = F_{01}^{II}.$$

The  $2 \times 2$  dimensionless electron collisional friction matrices,  $[F^{ee}](r)$ ,  $[F^{ei}](r)$ , and  $[F^{eI}](r)$ , are defined to have the following elements:  $F_{00}^{ee} = Z_{\text{eff}}$ ,  $F_{01}^{ee} = (3/2) Z_{\text{eff}}$ ,  $F_{10}^{ee} = F_{01}^{ee}$ ,  $F_{11}^{ee} = \sqrt{2} + (13/4) Z_{\text{eff}}$ ,  $F_{00}^{ei} = Z_{\text{eff}i}$ ,  $F_{01}^{ei} = F_{11}^{ei} = 0$ ,  $F_{10}^{ei} = (3/2) Z_{\text{eff}i}$ ,  $F_{00}^{eI} = Z_{\text{eff}I}$ ,  $F_{01}^{eI} = F_{11}^{eI} = 0$ ,  $F_{10}^{eI} = (3/2) Z_{\text{eff}I}$ .

### 2.4 Neoclassical Viscosity Matrices

The  $2 \times 2$  dimensionless species- $a$  neoclassical viscosity matrix,  $[\mu^a](r)$ , is defined to have the following elements:  $\mu_{00}^a = K_{00}^a$ ,  $\mu_{01}^a = (5/2) K_{00}^a - K_{01}^a$ ,  $\mu_{10}^a = \mu_{01}^a$ ,  $\mu_{11}^a = K_{11}^a - 5 K_{01}^a + (25/4) K_{00}^a$ . Here,

$$K_{jk}^e = g_t \frac{4}{3\sqrt{\pi}} \int_0^\infty \frac{e^{-x} x^{4+j+k} \nu_D^e(x) dx}{[x^2 + \nu_{*e} \nu_D^e(x)] [x^2 + (5\pi/8) (\omega_{te} \tau_{ee})^{-1} \nu_T^e(x)]}, \quad (53)$$

$$\nu_D^e = \frac{3\sqrt{\pi}}{4} \left[ \left(1 - \frac{1}{2x}\right) \psi(x) + \psi'(x) \right] + \frac{3\sqrt{\pi}}{4} Z_{\text{eff}}, \quad (54)$$

$$\nu_\epsilon^e = \frac{3\sqrt{\pi}}{2} [\psi(x) - \psi'(x)], \quad (55)$$

$$\nu_T^a(x) = 3\nu_D^a(x) + \nu_\epsilon^a(x), \quad (56)$$

and

$$\psi(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt - \frac{2}{\sqrt{\pi}} x e^{-x^2}, \quad (57)$$

$$\psi'(x) = \frac{2}{\sqrt{\pi}} x e^{-x^2}. \quad (58)$$

Furthermore,

$$K_{jk}^i = g_t \frac{4}{3\sqrt{\pi}} \int_0^\infty \frac{e^{-x} x^{2+j+k} \nu_D^i(x) dx}{[x + \nu_{*i} \nu_D^i(x)] [x + (5\pi/8) (\omega_{ti} \tau_{ii})^{-1} \nu_T^i(x)]}, \quad (59)$$

$$\begin{aligned} \nu_D^i &= \frac{3\sqrt{\pi}}{4} \left[ \left(1 - \frac{1}{2x}\right) \psi(x) + \psi'(x) \right] \frac{1}{x} \\ &+ \frac{3\sqrt{\pi}}{4} \alpha_I \left[ \left(1 - \frac{x_{iI}}{2x}\right) \psi\left(\frac{x}{x_{iI}}\right) + \psi'\left(\frac{x}{x_{iI}}\right) \right] \frac{1}{x}, \end{aligned} \quad (60)$$

$$\begin{aligned} \nu_\epsilon^i &= \frac{3\sqrt{\pi}}{2} [\psi(x) - \psi'(x)] \frac{1}{x} \\ &+ \frac{3\sqrt{\pi}}{2} \alpha_I \left[ \frac{m_i}{m_I} \psi\left(\frac{x}{x_{iI}}\right) - \psi'\left(\frac{x}{x_{iI}}\right) \right] \frac{1}{x}, \end{aligned} \quad (61)$$

and, finally,

$$K_{jk}^I = g_t \frac{4}{3\sqrt{\pi}} \int_0^\infty \frac{e^{-x} x^{2+j+k} \nu_D^I(x) dx}{[x + \nu_{*I} \nu_D^I(x)] [x + (5\pi/8) (\omega_{tI} \tau_{II})^{-1} \nu_T^I(x)]}, \quad (62)$$

$$\begin{aligned} \nu_D^I &= \frac{3\sqrt{\pi}}{4} \left[ \left(1 - \frac{1}{2x}\right) \psi(x) + \psi'(x) \right] \frac{1}{x} \\ &+ \frac{3\sqrt{\pi}}{4} \frac{1}{\alpha_I} \left[ \left(1 - \frac{x_{Ii}}{2x}\right) \psi\left(\frac{x}{x_{Ii}}\right) + \psi'\left(\frac{x}{x_{Ii}}\right) \right] \frac{1}{x}, \end{aligned} \quad (63)$$

$$\begin{aligned} \nu_\epsilon^I &= \frac{3\sqrt{\pi}}{2} [\psi(x) - \psi'(x)] \frac{1}{x} \\ &+ \frac{3\sqrt{\pi}}{2} \frac{1}{\alpha_I} \left[ \frac{m_I}{m_i} \psi\left(\frac{x}{x_{Ii}}\right) - \psi'\left(\frac{x}{x_{Ii}}\right) \right] \frac{1}{x}. \end{aligned} \quad (64)$$

Note that our expressions for the neoclassical viscosity matrices interpolate in the most accurate manner possible between the three standard neoclassical collisionality regimes (i.e., the banana, plateau, and Pfirsch-Schüter regimes).

## 2.5 Parallel Force and Heat Balance

Let  $[\tilde{\mu}^I] = \alpha_I^2 (T_i/T_I) x_{Ii} [\mu^I]$ . The requirement of equilibrium force and heat balance parallel to the magnetic field leads us to define four  $2 \times 2$  dimensionless ion matrices,  $[L^{ii}](r)$ ,  $[L^{iI}](r)$ ,



$[L^{Ii}](r)$ , and  $[L^{II}](r)$ , where

$$\begin{pmatrix} [L^{ii}], & [L^{iI}] \\ [L^{Ii}], & [L^{II}] \end{pmatrix} = \begin{pmatrix} [F^{ii} + \mu^i + Y^{in}/y_n], & -[F^{iI}] \\ -[F^{Ii}], & [F^{II} + \tilde{\mu}^I] \end{pmatrix}^{-1} \begin{pmatrix} [F^{ii} + Y^{in}], & -[F^{iI}] \\ -[F^{Ii}], & [F^{II}] \end{pmatrix}, \quad (65)$$

the additional four  $2 \times 2$  dimensionless ion matrices,  $[G^{ii}](r)$ ,  $[G^{iI}](r)$ ,  $[G^{Ii}](r)$ , and  $[G^{II}](r)$ , where

$$\begin{pmatrix} [G^{ii}], & [G^{iI}] \\ [G^{Ii}], & [G^{II}] \end{pmatrix} = \tau_{ii} \langle \sigma v \rangle_i^{\text{cx}} \langle n_n \rangle \begin{pmatrix} [F^{ii} + \mu^i + Y^{in}/y_n], & -[F^{iI}] \\ -[F^{Ii}], & [F^{II} + \tilde{\mu}^I] \end{pmatrix}^{-1}, \quad (66)$$

and the  $2 \times 2$  dimensionless electron matrices,  $[Q^{ee}](r)$ ,  $[G^{ei}](r)$ ,  $[L^{ee}](r)$ ,  $[L^{ei}](r)$ , and  $[L^{eI}](r)$ , where

$$[Q^{ee}] = [F^{ee} + \mu^e]^{-1}, \quad (67)$$

$$[G^{ei}] = [Q^{ee}] ([F^{ei}] [G^{ii}] + [F^{eI}] [G^{Ii}]), \quad (68)$$

$$[L^{ee}] = [Q^{ee}] [F^{ee}], \quad (69)$$

$$[L^{ei}] = [Q^{ee}] \{ [F^{ei}] [L^{ii}] - [F^{ei}] + [F^{eI}] [L^{Ii}] \}, \quad (70)$$

$$[L^{eI}] = [Q^{ee}] \{ [F^{eI}] [L^{II}] - [F^{eI}] + [F^{ei}] [L^{Ii}] \}. \quad (71)$$

Here,

$$[Y^{in}] = \tau_{ii} \langle \sigma v \rangle_i^{\text{cx}} \langle n_n \rangle \begin{bmatrix} 1, & 0 \\ 0, & E_n/T_i \end{bmatrix}, \quad (72)$$

$$y_n = \frac{\langle n_n \rangle \langle B^2 \rangle}{\langle n_n B^2 \rangle}, \quad (73)$$

where

$$\langle A \rangle(r) \equiv \oint \frac{A(r, \Theta) d\Theta}{B(r, \Theta)} \bigg/ \oint \frac{d\Theta}{B(r, \Theta)}. \quad (74)$$

Moreover,  $\langle \sigma v \rangle_i^{\text{cx}}$  is the flux-surface averaged rate constant for charge exchange reactions between neutrals and majority ions,  $n_n(r, \Theta)$  the neutral particle number density, and  $E_n/T_i$  the ratio of the incoming neutral energy to the majority ion energy. The parameter  $y_n$  takes into account the fact that the incoming neutrals at the edge of an H-mode tokamak plasma are usually concentrated at the X-point (i.e.,  $y_n > 1$ ).

## 2.6 Neoclassical Frequencies

The neoclassical frequencies of the three plasma species have the following definitions:

$$\omega_{\text{nc}i}(r) = -G_{00}^{ii} \omega_E - \left[ L_{00}^{ii} - L_{01}^{ii} \left( \frac{\eta_i}{1 + \eta_i} \right) \right] \omega_{*i} - \left[ L_{00}^{iI} - L_{01}^{iI} \left( \frac{\eta_I}{1 + \eta_I} \right) \right] \omega_{*I}, \quad (75)$$

$$\omega_{\text{nc}I}(r) = -G_{00}^{Ii} \omega_E - \left[ L_{00}^{II} - L_{01}^{II} \left( \frac{\eta_I}{1 + \eta_I} \right) \right] \omega_{*I} - \left[ L_{00}^{Ii} - L_{01}^{Ii} \left( \frac{\eta_i}{1 + \eta_i} \right) \right] \omega_{*i}, \quad (76)$$

$$\begin{aligned} \omega_{\text{nc}e}(r) = & -G_{00}^{ei} \omega_E - \left[ L_{00}^{ee} - L_{01}^{ee} \left( \frac{\eta_e}{1 + \eta_e} \right) \right] \omega_{*e} - \left[ L_{00}^{ei} - L_{01}^{ei} \left( \frac{\eta_i}{1 + \eta_i} \right) \right] \omega_{*i} \\ & - \left[ L_{00}^{eI} - L_{01}^{eI} \left( \frac{\eta_I}{1 + \eta_I} \right) \right] \omega_{*I}. \end{aligned} \quad (77)$$

Here,

$$\omega_E(r) = -\frac{d\Phi}{d\Psi_p}, \quad (78)$$

$$\omega_{*a}(r) = -\frac{T_a}{e_a} \frac{d \ln p_a}{d\Psi_p}, \quad (79)$$

$$\eta_a(r) = \frac{d \ln T_a}{d \ln n_a}. \quad (80)$$

Moreover,  $p_a(r) = n_a T_a$ , and  $\Phi(r)$  is the equilibrium electric scalar potential.

## 2.7 Impurity Ion Angular Rotation Velocities

Let

$$\omega_{\theta I}(r) = \frac{\mathbf{V}^I \cdot \nabla \theta}{\mathbf{B} \cdot \nabla \theta} \frac{R_0 B_0 g}{R^2}, \quad (81)$$

$$\omega_{\phi I}(r) = \mathbf{V}^I \cdot \nabla \phi, \quad (82)$$

where  $\mathbf{V}^I$  is the impurity ion fluid velocity, and the right-hand sides are evaluated on the outboard mid-plane. According to neoclassical theory,

$$\omega_{\theta I} = K_\theta \omega_{\text{nc}I}, \quad (83)$$

$$\omega_{\phi I} = \omega_E + \omega_{*I} + \omega_{\theta I}, \quad (84)$$

where

$$K_\theta(r) = \frac{R_0^2 B_0^2 g^2}{R^2 \langle B^2 \rangle}. \quad (85)$$

## 3 Plasma Response in Inner Region

### 3.1 Linear Layer Widths

Let

$$\tau_R(r) = \mu_0 r^2 \sigma_{ee} Q_{00}^{ee}, \quad (86)$$

$$\sigma_{ee}(r) = \frac{n_e e^2 \tau_{ee}}{m_e}, \quad (87)$$

$$\tau_H(r) = \frac{R_0}{B_0 g} \frac{\sqrt{\mu_0 \rho}}{n s}, \quad (88)$$

$$\rho(r) = m_i [n_i(r) + n_b(r)] + m_I n_I(r), \quad (89)$$

$$\tau_E(r) = \frac{r^2}{D_\perp + (2/3) \chi_e}, \quad (90)$$

$$\tau_M(r) = \frac{R_0^2 q^2}{\chi_\phi}, \quad (91)$$

$$\rho_s(r) = \frac{\sqrt{m_i T_e}}{e B_0 g}, \quad (92)$$

$$\tau(r) = -\frac{\omega_{*i}}{\omega_{*e}}, \quad (93)$$

$$S(r) = \frac{\tau_R}{\tau_H}, \quad (94)$$

$$P_M(r) = \frac{\tau_R}{\tau_M}, \quad (95)$$

$$P_E(r) = \frac{\tau_R}{\tau_E}, \quad (96)$$

$$D(r) = \frac{5}{3} S^{1/3} \frac{\rho_s}{r}, \quad (97)$$

$$Q_E(r) = S^{1/3} n \omega_E \tau_H, \quad (98)$$

$$Q_{e,i}(r) = -S^{1/3} n \omega_{*e,i} \tau_H \quad (99)$$

Here,  $\chi_e(r)$ ,  $D_\perp(r)$ , and  $\chi_\phi(r)$  are the perpendicular electron energy, particle, and toroidal momentum diffusivity profiles, respectively. Moreover,  $n_b(r)$  is the number density of fast (as opposed to thermal) majority ions. The constant- $\psi$  linear layer width is determined from the solution of

$$\frac{d^2 Y}{dp^2} - \left[ \frac{-Q_E (Q_E - Q_i) + i(Q_E - Q_i) (P_M + P_E) p^2 + P_M P_E p^4}{i(Q_E - Q_e) + \{P_E + i(Q_E - Q_i) D^2\} p^2 + (1 + \tau) P_M D^2 p^4} \right] p^2 Y = 0. \quad (100)$$

If the small- $p$  behavior of solution of the previous equation that is well-behaved as  $p \rightarrow \infty$  is

$$Y(p) = Y_0 [1 - c p + \mathcal{O}(p^2)]. \quad (101)$$

then the linear layer width is

$$\delta_{\text{linear}}(r) = \frac{\pi |c| r}{S^{1/3}}. \quad (102)$$

Note that Eq. (100) can only be solved when  $1 + \tau > 0$ .

### 3.2 Island Widths

Let  $\Psi_k = R_0 B_0 \hat{\Psi}_k e^{-i\varphi_k}$ , where  $\hat{\Psi}_k > 0$  and  $\varphi_k$  are both real quantities. The full width (in  $r$ ) of the magnetic island chain at the  $k$ th resonant surface is

$$W_k = 4 R_0 \left( \frac{q}{g s} \right)_{r_k}^{1/2} \hat{\Psi}_k^{1/2}. \quad (103)$$

### 3.3 Critical Island Widths

The critical full island width (in  $r$ ) which must be exceeded before the electron temperature is flattened within the magnetic separatrix of the magnetic island chain at the  $k$ th resonant surface is

$$W_{T_e k} = 5.07 \left( \frac{\chi_e}{\chi_{\parallel e}} \right)_{r_k}^{1/4} \left( \frac{1}{\epsilon s n} \right)_{r_k}^{1/2} r_k, \quad (104)$$

where  $\epsilon = r/R_0$ , and

$$\chi_{\parallel e} = \frac{\chi_{\parallel e}^{\text{brag}} \chi_{\parallel e}^{\text{max}}}{\chi_{\parallel e}^{\text{brag}} + \chi_{\parallel e}^{\text{max}}}, \quad (105)$$

$$\chi_{\parallel e}^{\text{brag}} = \frac{1.55 \tau_{ee} v_{Te}^2}{1 + 0.38 Z_{\text{eff}}}, \quad (106)$$

$$\chi_{\parallel e}^{\text{max}} = \frac{2 R_0 v_{Te}}{n s} \frac{r_k}{W_{T_e k}} \quad (107)$$

Equations (104)–(107) must be solved iteratively for  $W_{T_e k}/r_k$ .

The critical full island width (in  $r$ ) which must be exceeded before the ion temperature is flattened within the magnetic separatrix of the magnetic island chain at the  $k$ th resonant surface is

$$W_{T_i k} = 5.07 \left( \frac{\chi_i}{\chi_{\parallel i}} \right)_{r_k}^{1/4} \left( \frac{1}{\epsilon s n} \right)_{r_k}^{1/2} r_k, \quad (108)$$

where  $\chi_i(r)$  is the perpendicular ion energy diffusivity profile,

$$\chi_{\parallel i} = \frac{\chi_{\parallel i}^{\text{brag}} \chi_{\parallel i}^{\text{max}}}{\chi_{\parallel i}^{\text{brag}} + \chi_{\parallel i}^{\text{max}}}, \quad (109)$$

$$\chi_{\parallel i}^{\text{brag}} = 1.95 \tau_{ii} v_{Ti}^2, \quad (110)$$

$$\chi_{\parallel i}^{\text{max}} = \frac{2 R_0 v_{Ti}}{n s} \frac{r_k}{W_{T_i k}}. \quad (111)$$

Equations (108)–(111) must be solved iteratively for  $W_{T_i k}/r_k$ .

The critical full island width (in  $r$ ) which must be exceeded before the electron density is flattened within the magnetic separatrix of the magnetic island chain at the  $k$ th resonant surface is

$$W_{n_e k} = 5.07 \left( \frac{D_{\perp}}{\chi_{\parallel i}} \right)_{r_k}^{1/4} \left( \frac{1}{\epsilon s n} \right)_{r_k}^{1/2} r_k, \quad (112)$$

where

$$\chi_{\parallel i} = \frac{\chi_{\parallel i}^{\text{brag}} \chi_{\parallel i}^{\text{max}}}{\chi_{\parallel i}^{\text{brag}} + \chi_{\parallel i}^{\text{max}}}, \quad (113)$$

$$\chi_{\parallel i}^{\text{brag}} = 1.95 \tau_{ii} v_{Ti}^2, \quad (114)$$

$$\chi_{\parallel i}^{\text{max}} = \frac{2 R_0 v_{Ti}}{n s} \frac{r_k}{W_{n_e k}}. \quad (115)$$

Equations (112)–(115) must be solved iteratively for  $W_{n_e k}/r_k$ .

### 3.4 Resonant Plasma Response Model

Let  $\chi_k = R_0 B_0 \hat{\chi}_k e^{-i\zeta_k}$ , and  $E_{kk'} = \hat{E}_{kk'} e^{-i\xi_{kk'}}$ , where  $\hat{\chi}_k > 0$ ,  $\zeta_k$ ,  $\hat{E}'_{kk'} > 0$ , and  $\xi_{kk'}$  are all real quantities. Furthermore, let  $X_k = \hat{\Psi}_k \cos \varphi_k$  and  $Y_k = \hat{\Psi}_k \sin \varphi_k$ . The resonant plasma response model at the  $k$ th resonant surface takes the form

$$\begin{aligned} (\hat{W}_k + \hat{\delta}_k) \mathcal{S}_k \left( \frac{dX_k}{d\hat{t}} + \hat{\omega}_k Y_k \right) &= f_k X_k + \sum_{k'=1, K} \hat{E}_{kk'} (\cos \xi_{kk'} X_{k'} - \sin \xi_{kk'} Y_{k'}) \\ &\quad + \Sigma_{kw} U_k + \hat{E}_{kk} \hat{\chi}_k \cos \zeta_k, \end{aligned} \quad (116)$$

$$\begin{aligned} (\hat{W}_k + \hat{\delta}_k) \mathcal{S}_k \left( \frac{dY_k}{d\hat{t}} - \hat{\omega}_k X_k \right) &= f_k Y_k + \sum_{k'=1, K} \hat{E}_{kk'} (\cos \xi_{kk'} Y_{k'} + \sin \xi_{kk'} X_{k'}) \\ &\quad + \Sigma_{kw} V_k + \hat{E}_{kk} \hat{\chi}_k \sin \zeta_k, \end{aligned} \quad (117)$$

$$\hat{\tau}_w \frac{dU_k}{d\hat{t}} = \Delta_{kw} U_k + \Sigma_{kw} X_k, \quad (118)$$

$$\hat{\tau}_w \frac{dV_k}{d\hat{t}} = \Delta_{kw} V_k + \Sigma_{kw} Y_k, \quad (119)$$

where

$$\hat{W}_k = \frac{\mathcal{I} W_k}{2 r_k} = \frac{2 \mathcal{I}}{\epsilon_{100} \hat{r}_k} \left( \frac{q}{g s} \right)_{r_k}^{1/2} (X_k^2 + Y_k^2)^{1/4}, \quad (120)$$

$$\hat{\delta}_k = \frac{\delta_{\text{linear}}(r_k)}{R_0 \epsilon_{100} \hat{r}_k}, \quad (121)$$

$$\mathcal{S}_k = \frac{\tau_R(r_k)}{\tau_A}, \quad (122)$$

$$\tau_A = \left[ \frac{\mu_0 \rho(0) r_{100}^2}{B_0^2} \right]^{1/2}, \quad (123)$$

$$\hat{\tau}_w = \frac{\tau_w}{\tau_A}, \quad (124)$$

$$f_k = f_{bk} + f_{ck} + f_{pk} + f_{wk} + f_{sk}, \quad (125)$$

Here,  $\mathcal{I} = 0.8227$ ,  $\epsilon_{100} = r_{100}/R_0$ ,  $\hat{r} = r/r_{100}$ ,  $\hat{r}_k = r_k/r_{100}$ , and  $\hat{t} = t/\tau_A$ . Note that the EPEC resonant plasma response model interpolates smoothly between the linear regime and the nonlinear Rutherford regime. The linear regime corresponds to  $\hat{W}_k \ll \hat{\delta}_k$ , whereas the nonlinear region corresponds to  $\hat{W}_k \gg \hat{\delta}_k$ .

The perturbed bootstrap current terms in the response model take the form

$$f_{bk} = f_{bek} + f_{bik}, \quad (126)$$

$$f_{bek} = \alpha_{bek} (f_{bT_ek} + f_{bn_ek}), \quad (127)$$

$$f_{bik} = \alpha_{bik} (f_{bT_ik} + f_{bn_ik}), \quad (128)$$

$$f_{bT_ek} = \left( \frac{\eta_e}{1 + \eta_e} \right)_{r_k} \frac{\hat{W}_k}{\hat{W}_{T_ek}^2 + \hat{\rho}_{\theta ek}^2 + \hat{W}_k^2}, \quad (129)$$

$$f_{bn_ek} = \left( \frac{1}{1 + \eta_e} \right)_{r_k} \frac{\hat{W}_k}{\hat{W}_{n_ek}^2 + \hat{\rho}_{\theta ek}^2 + \hat{W}_k^2}, \quad (130)$$

$$f_{bT_ik} = \left( \frac{\eta_i}{1 + \eta_i} \right)_{r_k} \frac{\hat{W}_k}{\hat{W}_{T_ik}^2 + \hat{\rho}_{\theta ik}^2 + \hat{W}_k^2}, \quad (131)$$

$$f_{bn_ik} = \left( \frac{1}{1 + \eta_i} \right)_{r_k} \frac{\hat{W}_k}{\hat{W}_{n_ik}^2 + \hat{\rho}_{\theta ik}^2 + \hat{W}_k^2}, \quad (132)$$

where

$$\alpha_{bek} = -2\mathcal{I} I_g \left( \frac{\omega_{*e} + \omega_{nce}}{\omega_\beta} \right)_{r_k}, \quad (133)$$

$$\alpha_{bik} = 2\mathcal{I} I_g \left[ \frac{(n_i/n_e)(\omega_{*i} + \omega_{nci}) + (Z_I n_I/n_e)(\omega_{*I} + \omega_{ncI})}{\omega_\beta} \right]_{r_k}, \quad (134)$$

$$\omega_\beta(r) = \frac{sg B_0}{\mu_0 n_e e R_0^2 q}, \quad (135)$$

$$\hat{W}_{T_ek} = \frac{\mathcal{I} W_{T_ek}}{2 r_k}, \quad (136)$$

$$\hat{W}_{T_ik} = \frac{\mathcal{I} W_{T_ik}}{2 r_k}, \quad (137)$$

$$\hat{W}_{n_ek} = \frac{\mathcal{I} W_{n_ek}}{2 r_k}, \quad (138)$$

$$\hat{\rho}_{\theta e k} = \left( \frac{\mathcal{I} \rho_{\theta e}}{2r} \right)_{r_k} = \left( \frac{2\mathcal{I} v_{Te} m_e q R_0}{e B_0 g r^2} \right)_{r_k}, \quad (139)$$

$$\hat{\rho}_{\theta i k} = \left( \frac{\mathcal{I} \rho_{\theta i}}{2r} \right)_{r_k} = \left( \frac{2\mathcal{I} v_{Ti} m_i q R_0}{e B_0 g r^2} \right)_{r_k}, \quad (140)$$

and  $I_g = 1.58$ .

The curvature terms take the form

$$f_{ck} = \alpha_{ck} (f_{cT_e k} + f_{cn_e k} + f_{cT_i k} + f_{cn_i k}), \quad (141)$$

$$f_{cT_e k} = \left( \frac{n_e}{n_e + n_i} \frac{\eta_e}{1 + \eta_e} \right)_{r_k} \frac{\hat{W}_k}{\hat{W}_{T_e k}^2 + \hat{W}_k^2}, \quad (142)$$

$$f_{cn_e k} = \left( \frac{n_e}{n_e + n_i} \frac{1}{1 + \eta_e} \right)_{r_k} \frac{\hat{W}_k}{\hat{W}_{n_e k}^2 + \hat{W}_k^2}, \quad (143)$$

$$f_{cT_i k} = \left( \frac{n_i}{n_e + n_i} \frac{\eta_i}{1 + \eta_i} \right)_{r_k} \frac{\hat{W}_k}{\hat{W}_{T_i k}^2 + \hat{W}_k^2}, \quad (144)$$

$$f_{cn_i k} = \left( \frac{n_i}{n_e + n_i} \frac{1}{1 + \eta_i} \right)_{r_k} \frac{\hat{W}_k}{\hat{W}_{n_e k}^2 + \hat{W}_k^2}, \quad (145)$$

where

$$\alpha_{ck} = 2\mathcal{I} I_g D_R(r_k). \quad (146)$$

Here,  $D_R(r)$  is the Glasser-Greene-Johnson resistive interchange stability parameter.

The ion polarization terms take the form

$$f_{pk} = \alpha_{pk} (f_{pT_i k} + f_{pn_i k}), \quad (147)$$

$$f_{pT_i k} = \left( \frac{\eta_i}{1 + \eta_i} \right)_{r_k} \frac{\hat{W}_k}{(\hat{W}_{T_i k}^2 + \hat{W}_k^2)^2}, \quad (148)$$

$$f_{pn_i k} = \left( \frac{1}{1 + \eta_i} \right)_{r_k} \frac{\hat{W}_k}{(\hat{W}_{n_e k}^2 + \hat{W}_k^2)^2}, \quad (149)$$

where

$$\alpha_{pk} = 8\mathcal{I}^3 I_p \left[ \frac{(\omega_{*i} + \omega_{nci}) \omega_{nci}}{\omega_\beta \omega_\Omega} \right]_{r_k}, \quad (150)$$

$$\omega_\Omega(r) = \frac{e g B_0 s q}{m_i} \quad (151)$$

and  $I_p = 1.38$ .

The wall stabilization terms take the form

$$f_{wk} = \frac{\Sigma_{kw}^2}{\Delta_{kw}}. \quad (152)$$

Finally, the island saturation terms take the form

$$f_{sk} = - \left( \frac{f_{s1k} \hat{W}_{Te k}^2 + f_{s2k} \hat{W}_k^2}{\hat{W}_{Te k}^2 + \hat{W}_k^2} \right), \quad (153)$$

$$f_{s1k} = P_{1k} \frac{W_k}{r_k} \ln \left( \frac{r_k}{W_k} \right) + P_{2k} \frac{W_k}{r_k}, \quad (154)$$

$$f_{s2k} = P_{3k} \frac{W_k}{r_k}, \quad (155)$$

$$P_{1k} = 0.41 A_k^2, \quad (156)$$

$$P_{2k} = 0.41 \left( A_k^2 \left[ \ln \left( \frac{1}{r_k} \right) + 4.85 \right] - \frac{\Sigma_k^{nw} A_k}{2} - B_k - 0.44 A_k \right), \quad (157)$$

$$P_{3k} = 0.8 A_k^2 - 0.27 B_k - 0.09 A_k. \quad (158)$$

## 4 Plasma Angular Velocity Evolution

### 4.1 Evolution Equations

The quantity  $\hat{\omega}_k$  that appears in Eqs. (116) and (117) evolves in time according to

$$\hat{\omega}_k(\hat{t}) = \hat{\omega}_{k0} - \sum_{k'=1, K}^{p=1, \infty} \frac{m_k}{m_{k'}} \frac{y_p(\hat{r}_k)}{y_p(\hat{r}_{k'})} \alpha_{k', p}(\hat{t}) - \sum_{k'=1, K}^{p=1, \infty} \frac{z_p(\hat{r}_k)}{z_p(\hat{r}_{k'})} \beta_{k', p}(\hat{t}), \quad (159)$$

Here,  $\hat{\omega}_{k0} = \overline{\omega}_{k0} \tau_A$ ,  $y_p(\hat{r}) = J_1(j_{1,p} \hat{r}) / \hat{r}$ , and  $z_p(\hat{r}) = J_0(j_{0,p} \hat{r})$ . Moreover,  $\overline{\omega}_{k0}$  is the so-called ‘‘natural frequency’’ (in the absence of the RMP) at the  $k$ th resonant surface; this quantity is defined as the helical phase velocity of a naturally unstable island chain, resonant at the surface, in the absence of an RMP (or any other island chains). Furthermore,  $J_m(z)$  is a standard Bessel function, and  $j_{m,p}$  denotes the  $p$ th zero of this function. The time evolution equations for the  $\alpha_{k,p}$  and  $\beta_{k,p}$  parameters specify how the plasma poloidal and toroidal angular velocity profiles are modified by the electromagnetic torques that develop within the plasma, in response to the applied RMP, and how these modifications affect the natural frequencies. The evolution equations take the form

$$(1 + 2 Q_k^2) \frac{d\alpha_{k,p}}{d\hat{t}} + \left( \frac{j_{1,p}^2}{\hat{r}_{Mk}} + \frac{1}{\hat{r}_{\theta k}} + \frac{1}{\hat{r}_{cxk}} \right) \alpha_{k,p} = \frac{m_k^2 [y_p(\hat{r}_k)]^2}{\hat{\rho}_k \epsilon_{100}^2 [J_2(j_{1,p})]^2} \delta \hat{T}_k, \quad (160)$$

$$\frac{d\beta_{k,p}}{d\hat{t}} + \left( \frac{j_{0,p}^2}{\hat{r}_{Mk}} + \frac{1}{\hat{r}_{cxk}} \right) \beta_{k,p} = \frac{n^2 [z_p(\hat{r}_k)]^2}{\hat{\rho}_k [J_1(j_{0,p})]^2} \delta \hat{T}_k, \quad (161)$$



where

$$\begin{aligned} \delta \hat{T}_k &= \sum_{k'=1, K} \hat{E}_{kk'} [(Y_k X_{k'} - X_k Y_{k'}) \cos \xi_{kk'} - (X_k X_{k'} + Y_k Y_{k'}) \sin \xi_{kk'}] \\ &+ \Sigma_{kw} (Y_k U_k - X_k V_k) + \hat{E}_{kk} \hat{\chi}_k (Y_k \cos \zeta_k - X_k \sin \zeta_k). \end{aligned} \quad (162)$$

Here,  $Q_k = Q(r_k)$ ,  $\hat{\rho}_k = \rho(r_k)/\rho(0)$ ,  $\hat{\tau}_{Mk} = r_{100}^2/[\chi_\phi(r_k) \tau_A]$ ,  $\hat{\tau}_{\theta k} = \tau_\theta(r_k)/\tau_A$ ,  $\hat{\tau}_{\text{cx}k} = \tau_{\text{cx}}(r_k)/\tau_A$ . Moreover,

$$\tau_\theta(r) = \frac{\tau_{ii}}{\mu_{00}^i} \left/ \left( 1 + \frac{q^2 R_0^2}{r^2 a_{kk}} \right) \right. \quad (163)$$

is the poloidal flow damping timescale, and

$$\tau_{\text{cx}}(r) = \frac{1}{\langle n_n \rangle \langle \sigma v \rangle_i^{\text{cx}}} \quad (164)$$

is the charge exchange damping timescale. Furthermore,

$$Q^2(r) = \frac{q^2 R_0^2}{2r^2} \left( \left\langle \frac{1}{R^2} \right\rangle - \frac{1}{\langle R^2 \rangle} \right) \left/ \left\langle \frac{|\nabla r|^2}{R^2} \right\rangle \right. \quad (165)$$

## 4.2 Natural Frequencies

According to linear tearing mode theory, in the absence of the RMP, the natural frequency of the tearing mode resonant at the  $k$ th resonant surface is given by

$$\varpi_{ek} = -n (\omega_E + \omega_{*e})_{r_k}. \quad (166)$$

According to nonlinear tearing mode theory, in the absence of the RMP, the natural frequency of the tearing mode resonant at the  $k$ th resonant surface is given by

$$\varpi_{ik} = -n (\omega_E + \omega_{*i} + \omega_{\text{nc}i})_{r_k}. \quad (167)$$

It is also helpful to define

$$\varpi_{Ek} = -n (\omega_E)_{r_k}. \quad (168)$$

The EPEC model for the natural frequency is

$$\varpi_{k0} = \varpi_{Ek} + (\varpi_{ek} - \varpi_{Ek}) f_{ek} + (\varpi_{ik} - \varpi_{Ek}) f_{ik}, \quad (169)$$

where

$$f_{ek} = f_{Te k} + f_{ne k}, \quad (170)$$

$$f_{ik} = f_{Ti k} + f_{ni k}, \quad (171)$$

$$f_{Te k} = \left( \frac{\eta_e}{1 + \eta_e} \right)_{r_k} \frac{\hat{W}_{Te k}^2}{\hat{W}_{Te k}^2 + \hat{W}_k^2}, \quad (172)$$

$$f_{n_e k} = \left( \frac{1}{1 + \eta_e} \right)_{r_k} \frac{\hat{W}_{n_e k}^2}{\hat{W}_{n_e k}^2 + \hat{W}_k^2}, \quad (173)$$

$$f_{T_i k} = \left( \frac{\eta_i}{1 + \eta_i} \right)_{r_k} \frac{\hat{W}_k^2}{\hat{W}_{T_i k}^2 + \hat{W}_k^2}, \quad (174)$$

$$f_{n_i k} = \left( \frac{1}{1 + \eta_i} \right)_{r_k} \frac{\hat{W}_k^2}{\hat{W}_{n_e k}^2 + \hat{W}_k^2}. \quad (175)$$

## 5 Pressure Reduction

The full width (in  $\Psi_N$ ) of the magnetic separatrix of the island chain induced at the  $k$ th resonant surface is

$$\bar{W}_k = \left( \frac{d\Psi_N}{dr} \right)_{r_k} W_k = 4 \left( A_{1k} \hat{\Psi}_k \right)^{1/2}, \quad (176)$$

where

$$\frac{d\Psi_N}{dr} = \frac{r g}{R_0^2 q |\hat{\Psi}_p(0)|}, \quad (177)$$

$A_{1k} = A_1(r_k)$ ,  $A_1 = 1/(\sigma |\hat{\Psi}_p(0)|)$ ,  $\sigma(r) = d \ln q / d\Psi_N$ , and  $\hat{\Psi}_p(r) = \Psi_p / (R_0^2 B_0)$ . However, the halfway point between the inner and outer extremities of the separatrix is located at the flux-surface where  $\Psi_N = \Psi_N(r_k) - A_{2k} W_k^2 / 8$ ,  $A_{2k} = A_2(r_k)$ , and  $A_{2k} = [(1/3) (d^2 q / d\Psi_N^2) / (dq / d\Psi_N)]_{r_k}$ . On average, the presence of a magnetic island chain of full width  $W_k$  at the  $k$ th resonant surface causes the electron temperature, ion temperature, and electron number density profiles to be flattened in annular regions, centered on the halfway points between the inner and outer radii of the island separatrices, of widths (in  $\Psi_N$ )

$$\delta_{T_e k} = \frac{2}{\pi} \bar{W}_k \tanh \left( \frac{\bar{W}_k}{\bar{W}_{T_e k}} \right), \quad (178)$$

$$\delta_{T_i k} = \frac{2}{\pi} \bar{W}_k \tanh \left( \frac{\bar{W}_k}{\bar{W}_{T_i k}} \right), \quad (179)$$

$$\delta_{n_e k} = \frac{2}{\pi} \bar{W}_k \tanh \left( \frac{\bar{W}_k}{\bar{W}_{n_e k}} \right), \quad (180)$$

respectively, where

$$\bar{W}_{T_e k} = \left( \frac{d\Psi_N}{dr} \right)_{r_k} W_{T_e k}, \quad (181)$$

$$\bar{W}_{T_i k} = \left( \frac{d\Psi_N}{dr} \right)_{r_k} W_{T_i k}, \quad (182)$$

$$\bar{W}_{n_e k} = \left( \frac{d\Psi_N}{dr} \right)_{r_k} W_{n_e k}. \quad (183)$$

The total change in the plasma pressure interior to the  $j$ th resonant surface due to the temperature and density flattening at the various resonant surfaces (assuming that the pressure at the LCFS is fixed) is

$$\Delta P_j = \sum_{k=j,K} \left\{ (F_1 + F_3) \delta_{n_e k} + F_2 \delta_{T_e k} + F_4 \delta_{T_i k} + (F_5 + F_7) \delta_{n_e k} \bar{W}_k^2 + F_6 \delta_{T_e k} \bar{W}_k^2 + F_8 \delta_{T_i k} \bar{W}_k^2 + (F_9 + F_{11}) \delta_{n_e k}^3 + F_{10} \delta_{T_e k}^3 + F_{12} \delta_{T_i k}^3 \right\}_{r_k}, \quad (184)$$

where

$$F_1 = \frac{dn_e}{d\Psi_N} T_e, \quad (185)$$

$$F_2 = \frac{dT_e}{d\Psi_N} n_e, \quad (186)$$

$$F_3 = \frac{dn_i}{d\Psi_N} T_i, \quad (187)$$

$$F_4 = \frac{dT_i}{d\Psi_N} n_i, \quad (188)$$

$$F_5 = -\frac{A_2}{8} \left( \frac{dn_e}{d\Psi_N} \frac{dT_e}{d\Psi_N} + \frac{d^2 n_e}{d\Psi_n^2} T_e \right), \quad (189)$$

$$F_6 = -\frac{A_2}{8} \left( \frac{dn_e}{d\Psi_N} \frac{dT_e}{d\Psi_N} + \frac{d^2 T_e}{d\Psi_n^2} n_e \right), \quad (190)$$

$$F_7 = -\frac{A_2}{8} \left( \frac{dn_i}{d\Psi_N} \frac{dT_i}{d\Psi_N} + \frac{d^2 n_i}{d\Psi_n^2} T_i \right), \quad (191)$$

$$F_8 = -\frac{A_2}{8} \left( \frac{dn_i}{d\Psi_N} \frac{dT_i}{d\Psi_N} + \frac{d^2 T_i}{d\Psi_n^2} n_i \right), \quad (192)$$

$$F_9 = \frac{T_e}{24} \frac{d^3 n_e}{d\Psi_N^3}, \quad (193)$$

$$F_{10} = \frac{n_e}{24} \frac{d^3 T_e}{d\Psi_N^3}, \quad (194)$$

$$F_{11} = \frac{T_i}{24} \frac{d^3 n_i}{d\Psi_N^3}, \quad (195)$$

$$F_{12} = \frac{n_i}{24} \frac{d^3 T_i}{d\Psi_N^3}, \quad (196)$$

and we have neglected the impurity ion pressure.